

Table S5 : Efficacy of AN12855 in an Chronic (BALB/c) model of infection

Organ	Treatment	Conc (mg/kg)	Mean \pm SEM Log ₁₀ CFU (Number of mice)				
			Days post infection-1	27	41	55	83
Lung	Untreated		2.0 \pm 0.15 (3)	7.4 \pm 0.07(5)	6.3 \pm 0.13(5)	6.1 \pm 0.27(5)	6.3 \pm 0.11(5)
Lung	INH	25			5.7 \pm 0.08(5)*	5.0 \pm 0.07(5)*	3.7 \pm 0.09(5)*
Lung	AN12855	5			6.2 \pm 0.12(5)	6.3 \pm 0.13(5)	6.4 \pm 0.14(6)
Lung	AN12855	25			6.1 \pm 0.10(5)	5.9 \pm 0.03(5)	5.4 \pm 0.07(6)*
Lung	AN12855	100			5.8 \pm 0.08(5)*	5.3 \pm 0.07(6)	4.5 \pm 0.05(6)*
Spleen	Untreated			5.5 \pm 0.09(5)	4.8 \pm 0.16(5)	4.9 \pm 0.17(5)	5.5 \pm 0.05(5)
Spleen	INH	25			3.4 \pm 0.04(5)*	2.4 \pm 0.08(5)*	1.7 \pm 0.00(3) ^{A*}
Spleen	AN12855	5			4.8 \pm 0.14(5)	5.1 \pm 0.09(5)	5.3 \pm 0.11(5)
Spleen	AN12855	25			3.8 \pm 0.10(5)*	4.0 \pm 0.10(5)*	3.5 \pm 0.19(5)
Spleen	AN12855	100			3.4 \pm 0.10(5)*	2.9 \pm 0.13(5)*	2.1 \pm 0.10(5) ^{B*}

*: P value < 0.05

A: Data for two animals with CFU reported as the limit of detection, i.e. 50 CFU

B: Data for one animal with CFU reported as the limit of detection, i.e. 50 CFU