



# PGE<sub>2</sub> accounts for bidirectional changes in alveolar macrophage self-renewal with aging and smoking

Loka R Penke<sup>1,\*</sup> , Jennifer M Speth<sup>1,\*</sup>, Christina Draijer<sup>1</sup>, Zbigniew Zaslona<sup>1</sup>, Judy Chen<sup>2,4</sup> , Peter Mancuso<sup>2,3</sup>, Christine M Freeman<sup>1,2,6</sup>, Jeffrey L Curtis<sup>1,2,7</sup> , Daniel R Goldstein<sup>2,4,5</sup> , Marc Peters-Golden<sup>1,2</sup>

**Alveolar macrophages (AMs) are resident immune cells of the lung that are critical for host defense. AMs are capable of proliferative renewal, yet their numbers are known to decrease with aging and increase with cigarette smoking. The mechanism by which AM proliferation is physiologically restrained, and whether dysregulation of this brake contributes to altered AM numbers in pathologic circumstances, however, remains unknown. Mice of advanced age exhibited diminished basal AM numbers and contained elevated PGE<sub>2</sub> levels in their bronchoalveolar lavage fluid (BALF) as compared with young mice. Exogenous PGE<sub>2</sub> inhibited AM proliferation in an E prostanoïd receptor 2 (EP2)-cyclic AMP-dependent manner. Furthermore, EP2 knockout (EP2 KO) mice exhibited elevated basal AM numbers, and their AMs resisted the ability of PGE<sub>2</sub> and aged BALF to inhibit proliferation. In contrast, increased numbers of AMs in mice exposed to cigarette smoking were associated with reduced PGE<sub>2</sub> levels in BALF and were further exaggerated in EP2 KO mice. Collectively, our findings demonstrate that PGE<sub>2</sub> functions as a tunable brake on AM numbers under physiologic and pathophysiological conditions.**

DOI [10.26508/lsa.202000800](https://doi.org/10.26508/lsa.202000800) | Received 29 May 2020 | Revised 13 August 2020 | Accepted 14 August 2020 | Published online 20 August 2020

## Introduction

Alveolar macrophages (AMs) are the primary resident innate immune cells of the pulmonary alveoli. AMs recognize and clear inhaled pathogens and particulates, catabolize surfactant, and orchestrate the initiation and resolution of inflammation, which are critical to maintain lung homeostasis. AMs differ from other tissue macrophages in numerous respects; these differences have been attributed, at least in part, to the unique microenvironment of the alveolar space in which they reside (Hussell & Bell, 2014).

Tissue macrophages were long considered to arise predominantly from circulating monocytes. However, over the last decade, it has become well-accepted that at steady state, many resident tissue macrophages, including AMs, arise from embryonic precursors and maintain their numbers via self-renewal (Tarling et al, 1987; Guilliams et al, 2013; Hashimoto et al, 2013). Proliferative renewal is especially important in AMs because of their remarkable longevity in the lung (Maus et al, 2006). However, the regulation of proliferation in resident tissue macrophages, including AMs, remains poorly understood. It is worth noting that Gata6, a transcription factor implicated in the renewal of peritoneal macrophages (Rosas et al, 2014), is not expressed in AMs (Draijer et al, 2019). Nevertheless, AMs clearly proliferate in response to colony-stimulating factors GM-CSF and M-CSF (Chen et al, 1988; Draijer et al, 2019), with mTORC1 (Deng et al, 2017) and the transcription factors Bhlhe40 and Bhlhe41 (Rauschmeier et al, 2019) being responsible for this process. MafB, another transcription factor that is necessary for maintenance of AM numbers (Sato-Nishiwaki et al, 2013), is also expressed at very low levels in AMs (Hamada et al, 2020).

Common host-intrinsic and environmental factors have been linked to alterations in AM numbers in vivo. For example, reduced AM numbers contribute to the impaired pulmonary host defense associated with advanced age (Higashimoto et al, 1993; Wong et al, 2017). Conversely, an increased number of AMs associated with cigarette smoking (CS) (Hornby & Kellington, 1990; Barbers et al, 1991; Suzuki et al, 2020) have been implicated in the development of chronic obstructive pulmonary disease. Although these observations clearly establish that AM self-renewal is a tunable phenomenon, a unifying mechanism for such bidirectional alterations in AM numbers is lacking. Moreover, independent of recognized mechanisms promoting proliferation, the potential role of endogenous brakes on this process remains uncertain. Here, we used murine models of both aging and CS to reveal the importance of endogenous PGE<sub>2</sub> synthesis and signaling via the E prostanoïd 2

<sup>1</sup>Division of Pulmonary and Critical Care Medicine, Department of Internal Medicine, University of Michigan Medical School, Ann Arbor, MI, USA <sup>2</sup>Graduate Program in Immunology, University of Michigan Medical School, Ann Arbor, MI, USA <sup>3</sup>Department of Nutritional Sciences, University of Michigan School of Public Health, Ann Arbor, MI, USA <sup>4</sup>Division of Cardiology, Department of Internal Medicine, University of Michigan Medical School, Ann Arbor, MI, USA <sup>5</sup>Institute of Gerontology, University of Michigan, Ann Arbor, MI, USA <sup>6</sup>Research Service, Veterans Affairs Ann Arbor Healthcare System, Ann Arbor, MI, USA <sup>7</sup>Medical Service, Veterans Affairs Ann Arbor Healthcare System, Ann Arbor, MI, USA

Correspondence: [petersm@umich.edu](mailto:petersm@umich.edu)

\*Loka R Penke and Jennifer M Speth contributed equally to this work

receptor (EP2) as a novel brake on AM proliferative self-renewal that is itself subject to bidirectional modulation during perturbations.

## Results

### AM proliferative capacity is reduced in aged mice in association with increased levels of PGE<sub>2</sub> within the lung

Aging reduces the functions (Arahamian et al, 2008; Boe et al, 2017; Li et al, 2017) and numbers (Zissel et al, 1999; Wong et al, 2017) of AMs in both mice and humans. We analyzed AM numbers in the bronchoalveolar lavage fluid (BALF) of 18–22-mo-old mice compared with 6–8-wk young controls and found a significant reduction in AMs with aging (Fig 1A). These data confirm previous findings in which AMs were enumerated from mouse lung digests rather than BALF (Wong et al, 2017).

Reduced AM numbers with aging could result from cell-intrinsic or extrinsic factors. To evaluate the role of intrinsic factors, we isolated AMs from both young and aged mice and measured their proliferation after stimulation with the important AM mitogen GM-CSF (Hashimoto et al, 2013). Compared with young AMs, the proliferative capacity of aged AMs in response to GM-CSF was significantly reduced (Fig 1B). In parallel, aged AMs exhibited a similar, significant reduction in the expression of the cell cycle gene *Cyclin B1* compared with young AMs (Fig 1C). As extrinsic factors may shape functional changes in AMs (McQuattie-Pimentel et al, 2019 Preprint), we next evaluated the contribution of microenvironmental alterations in the aged lung to AM proliferation. To test this, we measured AM proliferation in young AMs cultured with GM-CSF in the presence of cell-free BALF from either young or aged mice. Whereas AM proliferation was intact when cells were cultured with BALF from young mice, it was diminished when they were cultured with BALF from aged mice (Fig 1D). These data suggest the presence of both cell-intrinsic defects and suppressive factors in the alveolar microenvironment that contribute to impaired AM proliferation associated with aging.

We recently reported that AMs from aged mice exhibited increased eicosanoid gene expression (Wong et al, 2017). We therefore used lipidomics to obtain an unbiased evaluation of lipid mediator content in BALF from both young and aged mice. BALF from aged mice contained altered levels of a variety of lipid mediators (Fig 1E); these include increases in those derived from the arachidonate cyclooxygenase (PGE<sub>2</sub> and 12-hydroxyheptadecatrienoic acid [12-HHT]) and lipoxygenase (leukotriene E<sub>4</sub> and 12-hydroxyeicosatetraenoic acid [12-HETE]) metabolic pathways and decreases in the linoleic acid oxidation derivative 13-hydroxyoctadecadienoic acid [13-HODE]. Of these lipid mediators, the greatest increment in BALF from aged relative to young mice was observed for PGE<sub>2</sub>. We used ELISA to confirm a significant elevation in PGE<sub>2</sub> within the BALF of an independent and larger cohort of aged mice (Fig 1F). To determine whether AM numbers and PGE<sub>2</sub> levels within the lungs of individual mice were related, correlation analysis was performed in a separate cohort of both young and aged mice. Indeed, there was a moderate but significant inverse correlation between AM numbers and PGE<sub>2</sub> levels of individual mice (Fig 1G).


### CS exposure induces AM proliferation concurrent with decreased PGE<sub>2</sub> production

In both human smokers and animal models of CS exposure, increased AM numbers (Hornby & Kellington, 1990; Barbers et al, 1991; Suzuki et al, 2020) accompany and have been implicated in a variety of forms of lung pathology and dysfunction (Jayes et al, 2016; Hu et al, 2019; Suzuki et al, 2020). However, there is conflicting evidence as to whether this reflects proliferation of resident AMs or recruitment of circulating monocytes (Golde et al, 1974; Hornby & Kellington, 1990; Hodge et al, 2007; Perez-Rial et al, 2013). We therefore enumerated resident AMs and assessed their proliferation in vivo in response to CS exposure (2 h/d for 7 d). Consistent with some previous studies, the total number of lavaged cells was significantly increased in mice exposed to 7 d of CS as compared with control mice exposed to room air (Fig 2A). This increase in total lavaged cells was accompanied by a significant increase in the frequency of resident AMs as identified by two independent sets of markers (CD11b<sup>+</sup>CD11c<sup>+</sup> and MerTK<sup>+</sup>CD64<sup>+</sup>) (Fig 2B). There were no significant differences in the number of CD11b<sup>+</sup>CD11c<sup>+</sup>Ly6G<sup>+</sup> neutrophils or CD11b<sup>+</sup>CD11c<sup>+</sup>CD103<sup>+</sup> dendritic cells (Fig 2C), suggesting that these populations did not contribute to increased cell numbers in CS-exposed mice. There was a nonsignificant trend in numbers of CD11b<sup>+</sup>CD11c<sup>+</sup>Ly6C<sup>+</sup> recruited monocytes in CS-exposed mice (Fig 2C); this is unlikely to be meaningful, however, given their very low absolute numbers relative to numbers of resident AMs. To confirm that this observed increase in AMs was due to enhanced proliferation, we measured proliferation via 5-ethynyl-2'-deoxyuridine (EdU) incorporation within resident AMs during CS exposure. CD11c<sup>+</sup> AMs isolated from the BALF of CS-exposed mice displayed a significant increase in EdU incorporation compared with cells from air-exposed mice (Fig 2D). Taken together, the results in this model of CS exposure indicate that the increase in AM numbers is due to increased AM proliferation, rather than an influx of circulating inflammatory monocytes.

Alveolar epithelial cells comprise the alveolar surface, and like AMs (Balter et al, 1989), they have the capacity to synthesize PGE<sub>2</sub> (Chauncey et al, 1988). Of note, ex vivo PGE<sub>2</sub> production by AMs was reduced in human smokers compared with nonsmokers (Balter et al, 1989) and in vitro CS exposure reduced PGE<sub>2</sub> synthesis by cultured airway epithelial cells obtained by brushing of human subjects (Zhang et al, 2011). We therefore hypothesized that the increase in AM numbers and proliferation in smoked mice may be associated with decreased PGE<sub>2</sub> levels on the alveolar surface. Indeed, CS exposure resulted in a time-dependent decrease in PGE<sub>2</sub> levels within the lung over 7 d, and this decrease persisted with continuous CS exposure for as long as 8 wk (Fig 2E).

### PGE<sub>2</sub> inhibits mitogen-induced AM proliferation via EP2-cAMP signaling


The above data establish that PGE<sub>2</sub> bioavailability within the lung is inversely correlated with the diametrically opposed abnormalities of AM numbers and proliferation in two models of abnormal lung immune function: aging and smoking. We next directly assessed the in vitro effect of PGE<sub>2</sub> on AM proliferation and interrogated its operative signaling mechanisms. Treatment of naïve AMs with 1 μM



**Figure 1. Alveolar macrophage (AM) proliferation is decreased in aged mice in association with elevated levels of PGE<sub>2</sub>.** (A) Absolute numbers of AMs enumerated from BALF of young (6–8 wk old) and aged (18–22 wk old) mice (n = 20–22 mice). (B) Primary AMs isolated from young and aged mice were treated with mouse GM-CSF (10 ng/ml) and incubated for 5 d and subjected to CyQuant proliferation assay measuring total cellular DNA (n = 5 mice). (C) Expression of *Cyclin B1* mRNA measured by RT-PCR in young and aged primary AMs treated with GM-CSF for 48 h (n = 3 separate experiments). (D-left panel) Schematic depicting young and aged BALF swap experiment. (D-right panel) Proliferation of young AMs incubated for 5 d in the presence of young or aged BALF, or medium alone, with or without GM-CSF (n = 5–9 mice). (E) Heat map of lipids that were elevated in BALF from both young (n = 4 mice) and aged (n = 3 mice) mice. (F) Levels of PGE<sub>2</sub> from young or aged BALF quantified by ELISA (n = 17–18 mice). (G) Pearson’s correlation analysis between AM numbers and PGE<sub>2</sub> levels in matched BALF samples from individual mice (n = 18 mice; 10 young, 8 aged). (A, B, C, D, E) Results in (B, C, D) are values expressed relative to those in unstimulated young AMs, and all data represent the mean ± SEM of the values indicated by individual symbols; ns, non-significant, \*P < 0.05, \*\*P < 0.01, \*\*\*P < 0.001; t test (A, E) or one-way ANOVA followed by Sidak’s test for multiple comparisons (B, C, D).

PGE<sub>2</sub> markedly reduced the ability of both GM-CSF and M-CSF to elicit AM proliferation (Fig 3A) and to induce expression of proliferation-associated genes (Fig 3B). We previously reported that among the four G protein-coupled receptors for PGE<sub>2</sub>, EP2 is

responsible for mediating most of its inhibitory effects on AM functions (Aronoff et al, 2004; Medeiros et al, 2009). Treatment with the selective EP2 agonist butaprost mimicked the ability of PGE<sub>2</sub> to inhibit AM proliferation in response to both GM-CSF and M-CSF (Fig



**Figure 2. Alveolar macrophage (AM) proliferation is reduced in cigarette smoking (CS)-exposed mice concurrent with decreased PGE<sub>2</sub> levels in the lung.** (A) Absolute numbers of lavaged cells in BALF collected from mice exposed to air or CS for 7 d (n = 10 mice). (B) Absolute numbers of CD11b<sup>-</sup>CD11c<sup>+</sup> and MerTK<sup>+</sup>CD64<sup>+</sup> resident AMs isolated from lung digests of mice exposed to air or CS (n = 3–8 mice), as measured by flow cytometry. (C) Absolute numbers of CD11b<sup>-</sup>CD11c<sup>-</sup>Ly6G<sup>+</sup> neutrophils, CD11b<sup>-</sup>CD11c<sup>-</sup>Ly6C<sup>+</sup> monocytes, and CD11b<sup>-</sup>CD11c<sup>-</sup>CD103<sup>+</sup> dendritic cells isolated from lung digests of mice exposed to air or CS (n = 3–8 mice), as measured by flow cytometry. (D) Frequency of EdU<sup>+</sup>CD11c<sup>+</sup> resident AMs isolated from the BALF of air or CS-exposed mice measured by flow cytometry. Mice were injected with EdU (1 mg/kg) 1 d before CS exposure and every other day for 7 total d (n = 9–10 mice). (E) Levels of PGE<sub>2</sub> from BALF of air or CS-exposed mice at baseline, 3 d, 5 d, 7 d and 8 wk (n = 3–5 mice). (A, B, C, D, E) Results are shown as mean ± SEM; \*P < 0.05, \*\*P < 0.01, \*\*\*P < 0.001; \*\*\*\*P < 0.0001; t test (A, D) or one-way ANOVA followed by Sidak's test for multiple comparisons (B, C, E).

3C). EP2 is a stimulatory G protein–coupled receptor that signals by activating adenylyl cyclase to increase intracellular production of cAMP (Aronoff et al, 2004). Forskolin, a direct activator of adenylyl cyclase, exerted similar inhibitory effects as PGE<sub>2</sub> and butaprost on mitogen-induced proliferation (Fig 3C).

The effects of cAMP signaling on AM functions are mediated by two distinct downstream cAMP effector molecules, PKA and guanine nucleotide exchange protein directly activated by cAMP-1 (Epac-1) (Cheng et al, 2008). Treatment with cAMP analogs that are selective agonists for either PKA (6-bnz-cAMP) or Epac-1 (8-pCPT-2'-O-Me-cAMP) both resulted in significant inhibition of AM proliferation in response to GM-CSF (Fig 3D). Collectively, these data suggest that the inhibitory effects of PGE<sub>2</sub> on AM proliferation proceed via an EP2-cAMP signaling pathway. Both cAMP effectors are capable of contributing to this suppressive effect.


#### Endogenous EP2 signaling is necessary for PGE<sub>2</sub> restraint of AM numbers and proliferation in vitro and in vivo

To validate that endogenous EP2 signaling restrains AM proliferative self-renewal, we used mice with a global deletion of the EP2

receptor (EP2 KO). The number of AMs lavaged from naïve EP2 KO mice was significantly higher than that from wild-type mice at baseline (Fig 4A). In addition, AMs from EP2 KO mice displayed an exaggerated proliferative capacity in response to GM-CSF stimulation (Fig 4B).

It was possible that the increased AM renewal in EP2 KO mice could reflect, at least in part, a stimulatory effect of PGE<sub>2</sub> via the now-unchecked actions of signaling via alternative PGE<sub>2</sub> receptors, particularly the known stimulatory receptors EP1 or 3. We therefore tested the direct effects of exogenous PGE<sub>2</sub> on proliferation of EP2 KO AMs in the presence or absence of GM-CSF. As expected, the ability of PGE<sub>2</sub> to inhibit AM proliferation in response to GM-CSF was lost in EP2 KO AMs (Fig 4C) confirming that PGE<sub>2</sub> acts via EP2 to attenuate this process. Moreover, PGE<sub>2</sub> had no effect—stimulatory or otherwise—on either basal or GM-CSF-stimulated proliferation of EP2 KO AMs (Fig 4C). These data indicate that the increased AM renewal in EP2 KO mice is explained entirely by removal of the PGE<sub>2</sub>-EP2 brake, rather than by any stimulatory actions of PGE<sub>2</sub> mediated via other EP receptors.

We next sought to determine the role of PGE<sub>2</sub>/EP2 signaling on abnormal AM proliferation and cell numbers demonstrated in the



**Figure 3. PGE<sub>2</sub> inhibits alveolar macrophage (AM) proliferation in response to mitogens via EP2-cAMP signaling.**

(A) Primary mouse AMs were cultured in the presence of M-CSF or GM-CSF (10 ng/ml) with or without PGE<sub>2</sub> (1 μM), and their proliferation was determined by CyQuant assay measuring total cellular DNA after 5 d (n = 6 mice). (B) Expression of *Cyclin D1* mRNA measured by RT-PCR in primary AMs treated with M-CSF or GM-CSF with or without PGE<sub>2</sub> for 48 h (n = 3 separate experiments). (C) Proliferation of primary AMs cultured for 5 d in the presence of either M-CSF or GM-CSF in the presence of forskolin (100 μM) or the EP2 agonist butaprost (1 μM) for over 5 d. \* = P < 0.05 compared with untreated control; § = P < 0.05 compared with forskolin alone; # = P < 0.05 compared with butaprost alone (n = 6 mice). (D) Proliferation of primary AMs cultured for 5 d in the presence of either M-CSF or GM-CSF in the presence of agonists for PKA (6-bnz-cAMP; 100 μM) and Epac-1 (8-pCPT-2'-O-Me-cAMP; 100 μM) (n = 4 mice). (A, B, C, D) Results are expressed relative to values of untreated control cells, and are shown as mean ± SEM; \*P < 0.05, \*\*P < 0.01; one-way ANOVA followed by Sidak's test for multiple comparisons (A, B, C, D).

mouse models of both aging and CS. We compared the effects of BALF from aged and young wild-type (WT) mice on proliferation of young AMs from both WT and EP2 KO mice. The ability of BALF from aged mice to suppress proliferation of AMs from WT mice was lost when the responder AMs were from EP2 KO mice (Fig 4D). This finding indicates that the effect of the suppressive factor within aged BALF—shown to contain increased levels of PGE<sub>2</sub> (Fig 1E and F)—depends on EP2 signaling within AMs and thus is unequivocally identified as PGE<sub>2</sub>. Furthermore, the increase in AMs within lung digests in response to CS exposure was further exaggerated in EP2 KO mice relative to WT mice (Fig 4E), in the absence of any significant increase in neutrophils, recruited monocytes, or dendritic cells (Fig 4F). The increase in AMs in mice exposed to CS has been linked with increased GM-CSF expression in the lung (Suzuki et al, 2020). By reducing lung PGE<sub>2</sub> concentration (Fig 2D), smoke exposure disrupts the physiologic brake on AM proliferative self-renewal, thereby mimicking the ability of EP2 deletion (Fig 4E) to yield exaggerated AM proliferation in response to mitogenic stimuli.


## Discussion

Studies investigating the mechanisms regulating AM proliferation are limited, and potential brakes on AM proliferation are poorly understood. One recent study implicated a suppressive role for

vitamin D<sub>3</sub> in GM-CSF-induced AM proliferation (Hu et al, 2019) that appears to play a role in the development of emphysema. Our studies revealed PGE<sub>2</sub> to be a novel endogenous brake on AM proliferation. Bidirectional perturbations in the lung levels of this lipid mediator contributed to alterations in AM-proliferative self-renewal in two clinically relevant models of lung immune dysfunction—aging and CS exposure.

In the instance of aging, PGE<sub>2</sub> levels were increased and correlated with decreased numbers of AMs within the lungs of mice of advanced age. This ability of PGE<sub>2</sub> to restrain AM self-renewal is consistent with its ability to inhibit proliferation of a broad variety of other cell types, including both immune (Gualde & Goodwin, 1982; Maric et al, 2018) and nonimmune (Moore et al, 2003; Huang et al, 2007; Michael et al, 2019) cells. Although PGE<sub>2</sub> has not, to our knowledge, been previously reported to inhibit proliferation of AMs or of any tissue macrophage population, such an effect is consistent with its ability to inhibit virtually all activation parameters of AMs, including phagocytosis, microbial killing, and inflammatory cytokine production (Aronoff et al, 2004, 2012; Serezani et al, 2007; Bourdonnay et al, 2012). Excess PGE<sub>2</sub> production by macrophages has been implicated in immune senescence (Wu & Meydani, 2014), although not specifically in the lung.

In contrast to the findings in aged mice, CS exposure resulted in significantly greater AM numbers in association with decreased levels of PGE<sub>2</sub> within the lung. The decline in PGE<sub>2</sub> levels was manifested within several days, reached their maximum by 7 d, and



**Figure 4. Loss of EP2 signaling enhances alveolar macrophage (AM) proliferation.**

(A) Absolute numbers of AMs isolated from the BALF of naïve WT or EP2 KO mice ( $n = 5$  mice). (B) Primary AMs were isolated from the BALF of WT and EP2 KO mice and cultured with GM-CSF (10 ng/ml) or medium alone for 5 d and then analyzed by CyQuant proliferation assay measuring total cellular DNA, using untreated WT as a control ( $n = 20$  mice). (C) Proliferation of primary EP2 KO AMs cultured for 5 d with PGE<sub>2</sub>, GM-CSF, or both. Values are expressed relative to untreated KO control cells. Dashed line represents WT baseline proliferation. ( $n = 6$  mice). (D) Proliferation of primary young WT or EP2 KO AMs cultured for 5 d with WT young or aged BALF in the presence or absence of GM-CSF ( $n = 3$  separate experiments; AMs from two individual mice per experiment). Values are expressed relative to untreated control for young or aged BALF,

persisted through 8 wk. The mechanism(s) by which CS exposure reduced BALF levels of PGE<sub>2</sub> was not addressed in our study. Whether CS targets AMs, alveolar epithelial cells or both cell types is uncertain. Moreover, PGE<sub>2</sub> levels reflect the balance of its synthesis, mediated by enzymes including cyclooxygenases and PGE synthases, and its degradation, mediated by 15-hydroxyprostaglandin dehydrogenase, and both classes of enzymes are subject to modulation (Tai et al, 2002; Stolberg et al, 2014). Finally, CS contains more than 5,000 discrete chemical compounds (Talhout et al, 2011), any of which, alone or in combination, could be responsible. Future research will be needed to explore the responsible compounds as well as target enzymes and cell types by which CS diminishes alveolar levels of PGE<sub>2</sub>. Likewise, the mechanism(s) responsible for the elevated PGE<sub>2</sub> levels in the alveolar space of aged mice remains to be determined. Notably, the consequences of excessive (aging) or impaired (smoking) PGE<sub>2</sub> production in the lung would be expected to extend beyond regulation of AM numbers and could contribute to global impairment and activation of AM functions in aging and smoking, respectively.

Both gain of function studies with the EP2-selective agonist butaprost and the direct adenylyl cyclase activator forskolin and loss-of-function studies employing cells from EP2 KO mice pointed to a critical role for EP2 signaling via cAMP in mediating the suppressive actions of PGE<sub>2</sub> on AM proliferation in vitro and in vivo. The similar magnitude of inhibition exerted by butaprost and forskolin suggests that EP2 agonism was sufficient to capture the full potential of cAMP-mediated inhibition of AM proliferation; this in turn argues against a significant role in this inhibition for the alternative stimulatory G protein-coupled PGE<sub>2</sub> receptor, EP4. Mechanistically, studies with PKA- and Epac-selective agonists revealed that both of these cAMP effectors were sufficient to account for inhibition of proliferation. In previous work, we have reported that roles for PKA and Epac-1 in regulating other AM functions can be either distinct or redundant (Aronoff et al, 2005), a phenomenon also demonstrated in other cell types (Cheng et al, 2008).

Our in vivo findings with EP2 KO mice suggest that endogenous levels of PGE<sub>2</sub> on the respiratory surface—which could derive from both AMs and alveolar epithelial cells—are sufficient to act via EP2 on AMs to restrain their numbers. Alternatively, our in vitro finding that EP2 KO AMs demonstrate enhanced ex vivo proliferation in response to GM-CSF can only be explained if AMs themselves are the source of PGE<sub>2</sub>, which in turn acts via EP2 as an autocrine brake on mitogen-stimulated proliferation. This, in turn, also suggests the possibility that the cell-intrinsic defect depicted in Fig 1B could also reflect excess autocrine production of PGE<sub>2</sub> by cultured AMs from aged mice. We speculate that a defect such as this that persists in culture may be mediated by epigenetic mechanisms; testing this possibility will require additional studies.

Although there are undoubtedly a number of determinants of AM self-renewal, our studies highlight the significance of PGE<sub>2</sub> as an endogenous suppressor of AM proliferation. Although we have

localized the PGE<sub>2</sub> effect to signaling via the EP2 receptor and downstream cAMP effectors, the molecular pathways involved in AM proliferation that are targeted for inhibition by PGE<sub>2</sub>-cAMP remain to be determined. However, it is interesting that PGE<sub>2</sub> is capable of inhibiting mTORC1—a critical driver of macrophage proliferation (Okunishi et al, 2014). Further studies will be necessary to determine the distinct mechanism(s) by which PGE<sub>2</sub> exerts its effects and whether alterations in its production or in EP2 expression influence AM self-renewal in other models of lung disease.

## Materials and Methods

### Animals

For isolation of naïve AMs, 6–8-wk-old WT C57Bl/6 female mice (Jackson Laboratory) were used. For CS experiments, 6–8-wk-old mice harboring a targeted deletion of both alleles of the *Ptger2* gene encoding the EP2 receptor (EP2 KO) were used. These mice were originally provided by Dr. Richard Breyer (Vanderbilt University) (Kennedy et al, 1999) and were maintained/bred by the University of Michigan Unit of Laboratory Medicine. For aging experiments, 8-wk-old (young) and 18–22-mo-old (aged) female C57Bl/6 mice were procured from Charles River or the National Institute of Aging Animal Facility, respectively. All mice were maintained at the University of Michigan Unit for Laboratory Animal Medicine. Animals were treated according to National Institutes of Health guidelines for the use of experimental animals with the approval of the University of Michigan Committee for the Use and Care of Animals.

### AM isolation and culture

Primary AMs were lavaged from mouse lungs as previously described (Bourdonnay et al, 2012). AMs were adhered for at least 1 h in RPMI with penicillin/streptomycin (pen/strep) and 2% FBS before treatment with appropriate compounds for indicated times.

### Quantitative real-time PCR

Total RNA was extracted from  $0.5 \times 10^6$  AMs using QIAGEN columns per manufacturer's instructions and converted to cDNA via reverse transcription. Relative gene expression of mouse cyclin B1 and cyclin D1 was determined by the  $\Delta$ Ct method, using SYBR Green dye (Applied Biosystems). Mouse  $\beta$ -actin was used as a reference gene. Primer sequences for mouse *Cyclin B1*, *Cyclin D1*, and  $\beta$ -actin were 5'-GAACCAGAGGTGGAAGCTTGC-3' (f), 5'-AGATGTTTCCATCGGGCTTG-3' (r); 5'-AGTGCCTGCAGAAGGAGATT-3' (f), 5'-AGGAAGCGTCCAGGTAGTT-3' (r); and 5'-GACGGCCAGGTCATCACTAT-3' (f), 5'-GCACTGTGTTGGCATA-GAGG-3' (r), respectively.

---

in both WT and KO cells. (E) Absolute numbers of CD11b<sup>+</sup>CD11c<sup>+</sup> and MerTK<sup>+</sup>CD64<sup>+</sup> resident AMs isolated from lung digests of EP2 KO mice exposed to air or cigarette smoking (n = 3–8 mice) measured by flow cytometry. Dashed line represents the average absolute number of all WT resident AMs for comparison. \*P < 0.05 compared with air control. (F) Absolute numbers of CD11b<sup>+</sup>Ly6G<sup>+</sup> neutrophils, CD11b<sup>+</sup>Ly6C<sup>+</sup> monocytes, and CD11c<sup>+</sup>CD103<sup>+</sup> dendritic cells isolated from lung digests of EP2 KO mice exposed to air or cigarette smoking (n = 3–8 mice) measured by flow cytometry. (A, B, C, D, E) Results are shown as mean  $\pm$  SEM; ns, nonsignificant, \*P < 0.05, \*\*P < 0.01; t test (A) one-way ANOVA followed by Sidak's test for multiple comparisons (B, C, D, E).

## Flow cytometric analysis of resident AMs

Lungs from mice subjected to CS or air were digested with collagenase as described (Stolberg et al, 2014) and single cell suspensions were labeled with antibodies for CD11c-V450 (clone N418; eBioscience), CD11b-AlexaFluor 700 (clone M1/70; eBioscience), MerTK-FITC (clone 2B10C42; BioLegend), CD64-APC (clone X54-5/7.1; BioLegend), Ly6C-PE/Cy7 (clone HK1.4; BioLegend), Ly6G-PE (clone 1A8; BioLegend), and CD103-PerCP/Cy5.5 (clone 2E7; BioLegend) at 4°C for 1 h. The population frequencies of resident AMs (CD11b<sup>+</sup>CD11c<sup>+</sup> or MerTK<sup>+</sup>CD64<sup>+</sup>), neutrophils (CD11b<sup>+</sup> Ly6G<sup>+</sup>), recruited monocytes (CD11c<sup>+</sup>CD11b<sup>+</sup>Ly6C<sup>+</sup>), and dendritic cells (CD11c<sup>+</sup>CD11b<sup>-</sup>CD103<sup>+</sup>) were analyzed on a BD Fortessa flow cytometer using BD FACSDiva software.

## Proliferation assay

Primary AMs were collected from mouse lung lavage fluid as described above. Cells were adhered in RPMI with 2% FBS and pen/strep in a 96-well plate at a concentration of 5,000 cells/well. Cells were treated with mitogen (mouse rM-CSF [10 ng/ml] or mouse rGM-CSF [10 ng/ml]; Peprotech) and incubated for 5 d. In some experiments, cells were pretreated with 1 μM PGE<sub>2</sub> (Cayman Chemical), 100 μM forskolin (Calbiochem), 1 μM EP2-selective agonist (butaprost; Cayman Chemical), 100 μM PKA agonist (6-bnz-cAMP; Axxora), or 100 μM Epac-1 agonist (8-pCPT-2'-O-Me-cAMP; Axxora) for 1 h before mitogen stimulation. Cell proliferation was determined after 5 d of culture using the CyQuant proliferation assay (Thermo Fisher Scientific) for DNA binding as per the manufacturer's instructions.

## Lipidomic analysis

BALF samples (800 μl) from young and aged mice (3–4 mice/group) were analyzed for lipid mediator content using an eicosanoid 44-plex lipidomic panel by Cayman Chemical. Briefly, BALF was subjected to liquid-phase and solid-phase extraction before immunoaffinity capture, then run through liquid chromatography and tandem mass spectrometry (LC instrument: Sciex ExionLC Integrated System, MS instrument: Sciex Triple Quad 6500+). Calculations of the total amount of each lipid present in the samples were performed using MultiQuant software (Sciex).

## PGE<sub>2</sub> ELISA

Individual mouse lungs were flushed with 800 μl PBS. The resulting lavage fluid was centrifuged at 500g for 10 min at 4°C to remove dead cells and debris, and PGE<sub>2</sub> levels were assessed by ELISA (Enzo Life Sciences).

## CS exposure

For in vivo CS exposure, mice were exposed to mainstream and side-stream CS as previously described (Phipps et al, 2010). Briefly, smoke from standardized 2R4F research cigarettes (University of Kentucky) was generated by a TE-2 CS machine (Teague Enterprises). Animals were exposed for 2 h/d and 7 d/wk in a whole-body exposure chamber. The mean concentration of particulates collected during a 2 h exposure was 19.05 ± 3.96 mg/m<sup>3</sup>/d. Control

animals were housed in an identical chamber but were exposed to room air with no smoke.

## EdU incorporation in CS-exposed AMs

For assessment of resident AM proliferation in response to CS exposure, mice were injected i.p. with 1 mg/kg EdU (BaseClick GmbH) immediately before CS exposure every other day for a total of four injections, starting 1 d before to the first CS exposure. Mice were euthanized within 2 h of final CS exposure and AMs were isolated from BALF as described previously (Bourdonnay et al, 2012). AMs from individual mice were labeled with fluorescently conjugated antibodies for CD11c and CD11b (sources as noted above), before fixation and permeabilization with BD CytoFix/CytoPerm reagent (BD Biosciences). EdU fluorescence was initiated using an in vivo EdU kit (BaseClick GmbH) per the manufacturer's instructions. CD11c<sup>+</sup>CD11b<sup>-</sup>EdU<sup>+</sup> AMs were enumerated via flow cytometry.

## Statistics

Data are presented as mean ± SEM unless otherwise specified. Differences between groups were tested using a one-way ANOVA followed by Sidak's test for multiple comparisons, or by a *t* test, as appropriate, using Prism 8.0 (GraphPad Software). For correlation between AM numbers and PGE<sub>2</sub> levels within the lungs of young and aged mice, Pearson's correlation analysis was performed using matched samples from 8 to 10 mice per group. *P*-values below 0.05 were considered to be significant.

## Supplementary Information

Supplementary Information is available at <https://doi.org/10.26508/lsa.202000800>.

## Acknowledgements

The authors would like to thank Rachel L Zemans for her assistance with the EdU incorporation assay procedure and members of the Peters-Golden laboratory for helpful input. The studies in this manuscript were supported by National Institutes of Health Grants R01 HL125555 and R35 HL144979 (to M Peters-Golden) and National Institutes of Health grants R01AG028082, R01HL120669, R01HL120669, R01AI138347, and K07AG050096 (to DR Goldstein). JM Speth and C Draijer were supported by National Institutes of Health T32 training grant HL 7749-23, JM Speth was also supported by Michigan Institute for Clinical & Health Research (MICHR) grant UL1TR00043 and American Cancer Society grant PF-17-143-01-TBG3. Z Zaslona was supported by an American Lung Association senior research fellowship award. P Mancuso was supported by Flight Attendants Medical Research Institute grant CIA-103071. J Chen is supported by National Institutes of Health T32 training grant T32AI007413. JL Curtis and CM Freeman were supported by Merit Review awards I01 CX000911 and I01 CX001553, respectively, from the Clinical Science Research & Development Service, Department of Veterans Affairs.

## Author Contributions

LR Penke: conceptualization, resources, data curation, formal analysis, validation, investigation, methodology, project administration, and writing—original draft, review, and editing.

JM Speth: conceptualization, resources, data curation, formal analysis, validation, investigation, methodology, project administration, and writing—original draft, review, and editing.

C Draijer: conceptualization, data curation, and validation.

Z Zaslona: conceptualization, data curation, and validation.

J Chen: data curation, validation, investigation, methodology, and writing—original draft.

P Mancuso: data curation, validation, investigation, and methodology.

CM Freeman: conceptualization, resources, data curation, and writing—original draft.

JL Curtis: conceptualization, data curation, investigation, and writing—original draft.

DR Goldstein: conceptualization, data curation, formal analysis, supervision, funding acquisition, validation, investigation, and writing—original draft.

M Peters-Golden: conceptualization, formal analysis, supervision, funding acquisition, validation, investigation, project administration, and writing—original draft, review, and editing.

## Conflict of Interest Statement

The authors declare that they have no conflict of interest.

## References

- Aprahamian T, Takemura Y, Goukassian D, Walsh K (2008) Ageing is associated with diminished apoptotic cell clearance in vivo. *Clin Exp Immunol* 152: 448–455. doi:10.1111/j.1365-2249.2008.03658.x
- Aronoff DM, Bergin IL, Lewis C, Goel D, O'Brien E, Peters-Golden M, Mancuso P (2012) E-prostanoid 2 receptor signaling suppresses lung innate immunity against *Streptococcus pneumoniae*. *Prostaglandins Other Lipid Mediat* 98: 23–30. doi:10.1016/j.prostaglandins.2012.03.002
- Aronoff DM, Canetti C, Peters-Golden M (2004) Prostaglandin E2 inhibits alveolar macrophage phagocytosis through an E-prostanoid 2 receptor-mediated increase in intracellular cyclic AMP. *J Immunol* 173: 559–565. doi:10.4049/jimmunol.173.1.559
- Aronoff DM, Canetti C, Serezani CH, Luo M, Peters-Golden M (2005) Cutting edge: Macrophage inhibition by cyclic AMP (cAMP): Differential roles of protein kinase A and exchange protein directly activated by cAMP-1. *J Immunol* 174: 595–599. doi:10.4049/jimmunol.174.2.595
- Balter MS, Toews GB, Peters-Golden M (1989) Multiple defects in arachidonate metabolism in alveolar macrophages from young asymptomatic smokers. *J Lab Clin Med* 114: 662–673.
- Barbers RG, Evans MJ, Gong H Jr., Tashkin DP (1991) Enhanced alveolar monocytic phagocyte (macrophage) proliferation in tobacco and marijuana smokers. *Am Rev Respir Dis* 143: 1092–1095. doi:10.1164/ajrccm/143.5\_pt\_1.1092
- Boe DM, Boule LA, Kovacs EJ (2017) Innate immune responses in the ageing lung. *Clin Exp Immunol* 187: 16–25. doi:10.1111/cei.12881
- Bourdonnay E, Serezani CH, Aronoff DM, Peters-Golden M (2012) Regulation of alveolar macrophage p40phox: Hierarchy of activating kinases and their inhibition by PGE<sub>2</sub>. *J Leukoc Biol* 92: 219–231. doi:10.1189/jlb.1211590
- Chauncey JB, Peters-Golden M, Simon RH (1988) Arachidonic acid metabolism by rat alveolar epithelial cells. *Lab Invest* 58: 133–140.
- Chen BD, Mueller M, Chou TH (1988) Role of granulocyte/macrophage colony-stimulating factor in the regulation of murine alveolar macrophage proliferation and differentiation. *J Immunol* 141: 139–144.
- Cheng X, Ji Z, Tsalkova T, Mei F (2008) Epac and PKA: A tale of two intracellular cAMP receptors. *Acta Biochim Biophys Sin (Shanghai)* 40: 651–662. doi:10.1111/j.1745-7270.2008.00438.x
- Deng W, Yang J, Lin X, Shin J, Gao J, Zhong XP (2017) Essential role of mTORC1 in self-renewal of murine alveolar macrophages. *J Immunol* 198: 492–504. doi:10.4049/jimmunol.1501845
- Draijer C, Penke LRK, Peters-Golden M (2019) Distinctive effects of GM-CSF and M-CSF on proliferation and polarization of two major pulmonary macrophage populations. *J Immunol* 202: 2700–2709. doi:10.4049/jimmunol.1801387
- Golde DW, Byers LA, Finley TN (1974) Proliferative capacity of human alveolar macrophage. *Nature* 247: 373–375. doi:10.1038/247373a0
- Gualde N, Goodwin JS (1982) Effects of prostaglandin E2 and preincubation on lectin-stimulated proliferation of human T cell subsets. *Cell Immunol* 70: 373–379. doi:10.1016/0008-8749(82)90339-2
- Guilliams M, De Kleer I, Henri S, Post S, Vanhoutte L, De Prijck S, Deswarte K, Malissen B, Hammad H, Lambrecht BN (2013) Alveolar macrophages develop from fetal monocytes that differentiate into long-lived cells in the first week of life via GM-CSF. *J Exp Med* 210: 1977–1992. doi:10.1084/jem.20131199
- Hamada M, Tsunakawa Y, Jeon H, Yadav MK, Takahashi S (2020) Role of MafB in macrophages. *Exp Anim* 69: 1–10. doi:10.1538/expanim.19-0076
- Hashimoto D, Chow A, Noizat C, Teo P, Beasley MB, Leboeuf M, Becker CD, See P, Price J, Lucas D, et al (2013) Tissue-resident macrophages self-maintain locally throughout adult life with minimal contribution from circulating monocytes. *Immunity* 38: 792–804. doi:10.1016/j.immuni.2013.04.004
- Higashimoto Y, Fukuchi Y, Shimada Y, Ishida K, Ohata M, Furuse T, Shu C, Teramoto S, Matsuse T, Sudo E, et al (1993) The effects of aging on the function of alveolar macrophages in mice. *Mech Ageing Dev* 69: 207–217. doi:10.1016/0047-6374(93)90024-I
- Hodge S, Hodge G, Ahern J, Jersmann H, Holmes M, Reynolds PN (2007) Smoking alters alveolar macrophage recognition and phagocytic ability: Implications in chronic obstructive pulmonary disease. *Am J Respir Cell Mol Biol* 37: 748–755. doi:10.1165/rcmb.2007-0025oc
- Hornby SB, Kellington JP (1990) DNA synthesis in alveolar macrophages and other changes in lavaged cells following exposure of CBA/H mice to cigarette smoke. *Environ Health Perspect* 85: 107–112. doi:10.2307/3430672
- Hu G, Dong T, Wang S, Jing H, Chen J (2019) Vitamin D3-vitamin D receptor axis suppresses pulmonary emphysema by maintaining alveolar macrophage homeostasis and function. *EBioMedicine* 45: 563–577. doi:10.1016/j.ebiom.2019.06.039
- Huang S, Wettlaufer SH, Hogaboam C, Aronoff DM, Peters-Golden M (2007) Prostaglandin E(2) inhibits collagen expression and proliferation in patient-derived normal lung fibroblasts via E prostanoid 2 receptor and cAMP signaling. *Am J Physiol Lung Cell Mol Physiol* 292: L405–L413. doi:10.1152/ajplung.00232.2006
- Hussell T, Bell TJ (2014) Alveolar macrophages: Plasticity in a tissue-specific context. *Nat Rev Immunol* 14: 81–93. doi:10.1038/nri3600
- Jays L, Haslam PL, Gratiou CG, Powell P, Britton J, Vardavas C, Jimenez-Ruiz C, Leonardi-Bee J, Tobacco Control Committee of the European Respiratory Society, (2016) SmokeHaz: Systematic reviews and meta-analyses of the effects of smoking on respiratory health. *Chest* 150: 164–179. doi:10.1016/j.chest.2016.03.060
- Kennedy CR, Zhang Y, Brandon S, Guan Y, Coffee K, Funk CD, Magnuson MA, Oates JA, Breyer MD, Breyer RM (1999) Salt-sensitive hypertension and reduced fertility in mice lacking the prostaglandin EP2 receptor. *Nat Med* 5: 217–220. doi:10.1038/5583
- Li Z, Jiao Y, Fan EK, Scott MJ, Li Y, Li S, Billiar TR, Wilson MA, Shi X, Fan J (2017) Aging-impaired filamentous actin polymerization signaling reduces alveolar macrophage phagocytosis of bacteria. *J Immunol* 199: 3176–3186. doi:10.4049/jimmunol.1700140

- Maric J, Ravindran A, Mazzurana L, Bjorklund AK, Van Acker A, Rao A, Friberg D, Dahlen SE, Heinemann A, Konya V, et al (2018) Prostaglandin E2 suppresses human group 2 innate lymphoid cell function. *J Allergy Clin Immunol* 141: 1761–1773.e6. doi:[10.1016/j.jaci.2017.09.050](https://doi.org/10.1016/j.jaci.2017.09.050)
- Maus UA, Janzen S, Wall G, Srivastava M, Blackwell TS, Christman JW, Seeger W, Welte T, Lohmeyer J (2006) Resident alveolar macrophages are replaced by recruited monocytes in response to endotoxin-induced lung inflammation. *Am J Respir Cell Mol Biol* 35: 227–235. doi:[10.1165/rcmb.2005-0241oc](https://doi.org/10.1165/rcmb.2005-0241oc)
- McQuattie-Pimentel AC, Ren Z, Joshi N, Watanabe S, Stoeger T, Chi M, Lu Z, Sichizya L, Piseaux R, Chen C-I, et al (2019) The aging microenvironment shapes alveolar macrophage identity in aging. *BioRxiv*. doi:[10.1101/717033](https://doi.org/10.1101/717033) Preprint posted July 29, 2019.
- Medeiros AI, Serezani CH, Lee SP, Peters-Golden M (2009) Efferocytosis impairs pulmonary macrophage and lung antibacterial function via PGE2/EP2 signaling. *J Exp Med* 206: 61–68. doi:[10.1084/jem.20082058](https://doi.org/10.1084/jem.20082058)
- Michael JV, Gavrila A, Nayak AP, Pera T, Liberato JR, Polischak SR, Shah SD, Deshpande DA, Penn RB (2019) Cooperativity of E-prostanoid receptor subtypes in regulating signaling and growth inhibition in human airway smooth muscle. *FASEB J* 33: 4780–4789. doi:[10.1096/fj.201801959r](https://doi.org/10.1096/fj.201801959r)
- Moore BB, Peters-Golden M, Christensen PJ, Lama V, Kuziel WA, Paine R 3rd, Toews GB (2003) Alveolar epithelial cell inhibition of fibroblast proliferation is regulated by MCP-1/CCR2 and mediated by PGE2. *Am J Physiol Lung Cell Mol Physiol* 284: L342–L349. doi:[10.1152/ajplung.00168.2002](https://doi.org/10.1152/ajplung.00168.2002)
- Okunishi K, DeGraaf AJ, Zaslona Z, Peters-Golden M (2014) Inhibition of protein translation as a novel mechanism for prostaglandin E2 regulation of cell functions. *FASEB J* 28: 56–66. doi:[10.1096/fj.13-231720](https://doi.org/10.1096/fj.13-231720)
- Perez-Rial S, del Puerto-Nevado L, Terron-Exposito R, Giron-Martinez A, Gonzalez-Mangado N, Peces-Barba G (2013) Role of recently migrated monocytes in cigarette smoke-induced lung inflammation in different strain of mice. *PLoS One* 8: e72975. doi:[10.1371/journal.pone.0072975](https://doi.org/10.1371/journal.pone.0072975)
- Phipps JC, Aronoff DM, Curtis JL, Goel D, O'Brien E, Mancuso P (2010) Cigarette smoke exposure impairs pulmonary bacterial clearance and alveolar macrophage complement-mediated phagocytosis of *Streptococcus pneumoniae*. *Infect Immun* 78: 1214–1220. doi:[10.1128/iai.00963-09](https://doi.org/10.1128/iai.00963-09)
- Rauschmeier R, Gustafsson C, Reinhardt A, Noelia A-G, Tortola L, Cansever D, Subramanian S, Taneja R, Rossner MJ, Sieweke MH, et al (2019) Bhlhe40 and Bhlhe41 transcription factors regulate alveolar macrophage self-renewal and identity. *EMBO J* 38: e101233. doi:[10.15252/embj.2018101233](https://doi.org/10.15252/embj.2018101233)
- Rosas M, Davies LC, Giles PJ, Liao CT, Kharfan B, Stone TC, O'Donnell VB, Fraser DJ, Jones SA, Taylor PR (2014) The transcription factor Gata6 links tissue macrophage phenotype and proliferative renewal. *Science* 344: 645–648. doi:[10.1126/science.1251414](https://doi.org/10.1126/science.1251414)
- Sato-Nishiwaki M, Aida Y, Abe S, Shibata Y, Kimura T, Yamauchi K, Kishi H, Igarashi A, Inoue S, Sato M, et al (2013) Reduced number and morphofunctional change of alveolar macrophages in MafB gene-targeted mice. *PLoS One* 8: e73963. doi:[10.1371/journal.pone.0073963](https://doi.org/10.1371/journal.pone.0073963)
- Serezani CH, Chung J, Ballinger MN, Moore BB, Aronoff DM, Peters-Golden M (2007) Prostaglandin E2 suppresses bacterial killing in alveolar macrophages by inhibiting NADPH oxidase. *Am J Respir Cell Mol Biol* 37: 562–570. doi:[10.1165/rcmb.2007-0153oc](https://doi.org/10.1165/rcmb.2007-0153oc)
- Stolberg VR, Martin B, Mancuso P, Olszewski MA, Freeman CM, Curtis JL, Chensue SW (2014) Role of CC chemokine receptor 4 in natural killer cell activation during acute cigarette smoke exposure. *Am J Pathol* 184: 454–463. doi:[10.1016/j.ajpath.2013.10.017](https://doi.org/10.1016/j.ajpath.2013.10.017)
- Suzuki T, McCarthy C, Carey BC, Borchers M, Beck D, Wikenheiser-Brokamp KA, Black D, Chalk C, Trapnell BC (2020) Increased pulmonary GM-CSF causes alveolar macrophage accumulation. Mechanistic implications for desquamative interstitial pneumonitis. *Am J Respir Cell Mol Biol* 62: 87–94. doi:[10.1165/rcmb.2018-0294oc](https://doi.org/10.1165/rcmb.2018-0294oc)
- Tai HH, Ensor CM, Tong M, Zhou H, Yan F (2002) Prostaglandin catabolizing enzymes. *Prostaglandins Other Lipid Mediat* 68-69: 483–493. doi:[10.1016/s0090-6980\(02\)00050-3](https://doi.org/10.1016/s0090-6980(02)00050-3)
- Talhout R, Schulz T, Florek E, van Benthem J, Wester P, Opperhuizen A (2011) Hazardous compounds in tobacco smoke. *Int J Environ Res Public Health* 8: 613–628. doi:[10.3390/ijerph8020613](https://doi.org/10.3390/ijerph8020613)
- Tarling JD, Lin HS, Hsu S (1987) Self-renewal of pulmonary alveolar macrophages: Evidence from radiation chimera studies. *J Leukoc Biol* 42: 443–446. doi:[10.1002/jlb.42.5.443](https://doi.org/10.1002/jlb.42.5.443)
- Wong CK, Smith CA, Sakamoto K, Kaminski N, Koff JL, Goldstein DR (2017) Aging impairs alveolar macrophage phagocytosis and increases influenza-induced mortality in mice. *J Immunol* 199: 1060–1068. doi:[10.4049/jimmunol.1700397](https://doi.org/10.4049/jimmunol.1700397)
- Wu D, Meydani SN (2014) Age-associated changes in immune function: Impact of vitamin E intervention and the underlying mechanisms. *Endocr Metab Immune Disord Drug Targets* 14: 283–289. doi:[10.2174/1871530314666140922143950](https://doi.org/10.2174/1871530314666140922143950)
- Zhang W, Case S, Bowler RP, Martin RJ, Jiang D, Chu HW (2011) Cigarette smoke modulates PGE(2) and host defence against *Moraxella catarrhalis* infection in human airway epithelial cells. *Respirology* 16: 508–516. doi:[10.1111/j.1440-1843.2010.01920.x](https://doi.org/10.1111/j.1440-1843.2010.01920.x)
- Zissel G, Schlaak M, Muller-Quernheim J (1999) Age-related decrease in accessory cell function of human alveolar macrophages. *J Investig Med* 47: 51–56.



**License:** This article is available under a Creative Commons License (Attribution 4.0 International, as described at <https://creativecommons.org/licenses/by/4.0/>).