



Correction



Life Science Alliance

Correction: Reduced protein-coding transcript diversity in severe dengue emphasises the role of alternative splicing

Priyanka Mehta^{1,2,*} , Chinky Shiu Chen Liu^{1,*}, Sristi Sinha¹, Ramakant Mohite¹, Smriti Arora¹, Partha Chattopadhyay^{1,2}, Sandeep Budhiraja³, Bansidhar Tarai³, Rajesh Pandey^{1,2} 

¹Division of Immunology and Infectious Disease Biology, Integrative GENomics of HOst-PathogEn (INGEN-HOPE) Laboratory, CSIR-Institute of Genomics and Integrative Biology (CSIR-IGIB), Delhi, India ²Academy of Scientific and Innovative Research (AcSIR), Ghaziabad, India ³Max Super Speciality Hospital (A Unit of Devki Devi Foundation), Max Healthcare, Delhi, India

Correspondence: rajesh.p@igib.res.in, rajeshp@igib.in

*Priyanka Mehta and Chinky Shiu Chen Liu contributed equally to this work

DOI <https://doi.org/10.26508/lsa.202402882> | Received 10 June 2024 | Accepted 11 June 2024 | Published online 14 June 2024

Article: Mehta P, Liu CSC, Sinha S, Mohite R, Arora S, Chattopadhyay P, Budhiraja S, Tarai B, Pandey R (2024 Jun 3) Reduced protein-coding transcript diversity in severe dengue emphasises the role of alternative splicing. *Life Sci Alliance* 7(8): e202402683. doi: [10.26508/lsa.202402683](https://doi.org/10.26508/lsa.202402683). PMID: 38830771.

Table 1. Clinical characteristics of the patient cohort.

Clinical parameters	Mild (n = 45)	Moderate (n = 46)	Severe (n = 21)	P-values
Age	27 (18–33)	26 (19–31.75)	35 (18–43)	0.14
Gender F M	15 30	17 29	4 17	0.34
NS1 antigen	3.22 (0.83–3.60)	3.5 (0.86–3.5)	3.5 (1.9–3.5)	0.75 ^a
Total leucocyte count (TLC)	5 (4.6–5.9)	3 (2.5–3.575)	2.7 (2.3–3.1)	<0.001
RBC count	4.76 (4.51–4.99)	4.835 (4.58–5.02)	4.78 (4.51–5.58)	0.76
Packed cell, volume	41.2 (40.1–42.8)	42.25 (38.4–44.22)	43.9 (40.4–46.3)	0.18
Platelet count	175 (155–202)	160 (150–165)	125 (112–135)	<0.001
Neutrophils	75.9 (66.7–79.5)	63.3 (58.1–68.65)	62.6 (57.3–73)	<0.001
Lymphocytes	14.1 (9.4–21.3)	24.5 (20.3–31.975)	26 (15.8–32.7)	<0.001
Monocytes	9.8 (8–12)	10 (7.85–11.475)	9 (6.9–11)	0.32
Eosinophils	0.2 (0–0.725)	0.2 (0.1–0.875)	0.5 (0.1–1)	0.32
Basophils	0.5 (0.3–0.6)	0.6 (0.225–0.8)	0.5 (0.3–0.8)	0.69
Hb	13.6 (12.75–14.3)	13.9 (12.52–14.5)	14.1 (13.1–15.3)	0.28
Total protein	7.3 (7–7.6)	6.7 (6.5–7.3)	6.7 (6.05–7.125)	0.03^a
Albumin	4.4 (4.05–4.5)	4 (3.8–4.15)	4 (3.675–4.1)	0.04^a
Globulin	3.1 (2.95–3.3)	3 (2.65–3.15)	2.65 (2.3–2.925)	0.06 ^a
A.G. ratio	1.4 (1.3–1.55)	1.3 (1.2–1.5)	1.55 (1.375–1.6)	0.19 ^a
Bilirubin (total)	0.7 (0.55–0.8)	0.4 (0.3–0.55)	0.75 (0.5–0.9)	0.02^a
Bilirubin (direct)	0.135 (0.12–0.17)	0.09 (0.07–0.13)	0.17 (0.13–0.23)	0.02^a
Bilirubin (indirect)	0.56 (0.43–0.615)	0.31 (0.24–0.42)	0.55 (0.39–0.71)	0.03^a
SGOT- aspartate transaminase (AST)	41 (34.5–76)	59 (44–80.5)	97.5 (74.25–141.5)	0.03^a

(Continued on following page)

Table 1. Continued

Clinical parameters	Mild (n = 45)	Moderate (n = 46)	Severe (n = 21)	P-values
SGPT- alanine transaminase (ALT)	42 (20–95)	41 (25.5–59)	66 (51.75–80.25)	0.13 ^a
AST/ALT ratio	0.93 (0.82–1.725)	1.48 (1.13–1.95)	1.705 (1.2–2.17)	0.08 ^a
Alkaline phosphatase	67 (60–89)	80 (59.5–100)	82.5 (59.75–124.25)	0.61 ^a
GGTP (gamma GT), serum	37 (24.5–46.5)	19 (14–41)	49 (32–82)	0.08 ^a

Data represented as median (IQR) or n (%).

^aIncomplete data points in either group.



License: This article is available under a Creative Commons License (Attribution 4.0 International, as described at <https://creativecommons.org/licenses/by/4.0/>).